TECHNICAL DATA SHEET · ANTIBACTERIALS

WELBMD100 – Bacitracin Methylene Disalicylate (BMD) 10% Premix

Polypeptide antibiotic premix for in-feed use, containing 100 mg bacitracin (as bacitracin methylene disalicylate) per gram (10%, 4,000 units per gram).

Pigs | Broiler chickens | Meat ducks | Mixed in feed | 25 kg/bag



TARGET ANIMALS

- Pigs
- Broiler chickens
- Meat ducks

WITHDRAWAL PERIOD

Pigs, chickens, ducks	0 days

INDICATIONS

- **Pigs, broiler chickens and meat ducks:** promoting growth.

Mode of action: Bacitracin inhibits bacterial cell-wall peptidoglycan synthesis. Good activity against most Gram-positive bacteria (Staphylococcus, Streptococcus, Enterococcus, Clostridium, Corynebacterium); also active against actinomycetes and spirochaetes. Resistance rarely develops. Synergistic with penicillin, streptomycin, neomycin and colistin; combinations with colistin are antagonistic with oxytetracycline, chlortetracycline, kitasamycin, enramycin, virginiamycin and olaquindox.

DOSAGE & ADMINISTRATION

Animal	Bacitracin per 1,000 kg feed	WELBMD100 per 1,000 kg feed	Duration
Pigs	10–30 g	100–300 g	—
Broiler chickens	5–40 g	50–400 g	—
Meat ducks	5–50 g	50–500 g	Not more than 42 days continuously

Administration: mixed in feed. Dosage expressed as bacitracin; product amounts calculated for WELBMD100 (100 mg/g).

ADVERSE REACTIONS

- Strong renal toxicity may occur after injection (this product is for in-feed use only).

PRECAUTIONS

- Do not use in breeding poultry.

Please use all products in accordance with local laws and regulations.

Need a quotation, COA or samples of WELBMD100?

Full product details on wellsunpharma.com · email anthony@wlsunshine.com

Request a Quote

This technical data sheet is for professional reference only. Specifications may be updated without notice; the COA supplied with each batch is the binding quality document. TDS-WELBMD100 · Rev. 1 · Issued November 2026.