TECHNICAL DATA SHEET · ANTICOCCIDIALS

WELCLOPI250 – Clopidol 25% Premix

Pyridinol anticoccidial premix for in-feed use, containing 250 mg clopidol per gram (25%).

Chickens | Rabbits | Mixed in feed | 25 kg/bag



TARGET ANIMALS

- Chickens
- Rabbits

WITHDRAWAL PERIOD

Chickens	5 days
Rabbits	5 days

INDICATIONS

- **Chickens and rabbits:** prevention of coccidiosis. Strongest action against *Eimeria tenella*; also active against rabbit coccidia.
- Suited to prevention only; it has no therapeutic value once coccidiosis has developed.

Mode of action: Clopidol mainly inhibits the sporozoite stage (peak activity on day 1 after infection), so it must be given before or at the time of infection.

DOSAGE & ADMINISTRATION

Animal	WELCLOPI250 per 1,000 kg feed	Clopidol per 1,000 kg feed
Chickens	500 g	125 g
Rabbits	800 g	200 g

Administration: mixed in feed. Dose expressed as the product; clopidol amounts calculated for 25%.

ADVERSE REACTIONS

- No adverse reactions have been observed at the recommended dosage.

PRECAUTIONS

- Can be used in commercial feed and during the rearing period.
- Do not use in laying hens producing eggs for human consumption during the laying period.
- Clopidol suppresses immunity to coccidia; withdrawing it too early can lead to coccidiosis outbreaks.
- Replacement pullets can be fed continuously up to 16 weeks of age.
- On farms with clopidol-resistant coccidia, do not switch to quinolone anticoccidials such as decoquinate.

Please use all products in accordance with local laws and regulations.

Need a quotation, COA or samples of WELCLOPI250?

Full product details on wellsunpharma.com · email anthony@wlsunshine.com

Request a Quote

This technical data sheet is for professional reference only. Specifications may be updated without notice; the COA supplied with each batch is the binding quality document. TDS-WELCLOPI250 · Rev. 1 · Issued November 2026.