TECHNICAL DATA SHEET · ANTIPARASITICS & FLY CONTROL

WELCYRO10 – Cyromazine 1% Premix

Insect growth regulator premix for fly control, containing 10 mg cyromazine per gram (1%).

Chickens | Mixed in feed | 25 kg/bag



TARGET ANIMALS

- Chickens

WITHDRAWAL PERIOD

Chickens	3 days

INDICATIONS

- **Fly control:** controls the breeding of fly larvae in the manure of animal houses.

Mode of action: Cyromazine stops dipteran larvae, especially first-stage larvae, from moulting; pupae develop abnormally and adult flies cannot emerge. Most is excreted unchanged in the droppings.

DOSAGE & ADMINISTRATION

Animal	WELCYRO10 per 1,000 kg feed	Cyromazine in feed	Duration
Chickens	500 g	5 mg/kg	4–6 weeks

Administration: mixed in feed. Dose expressed as the product.

ADVERSE REACTIONS

- No adverse reactions have been observed at the recommended dosage.

PRECAUTIONS

- At 25 mg/kg of feed feed intake may increase; at 500 mg/kg or more it falls; long-term feeding at 1,000 mg/kg or more may cause death from reduced intake.
- Apply manure from treated chickens to land at 1,000–2,000 kg per hectare; more than 9,000 kg per hectare may harm plant growth.

Please use all products in accordance with local laws and regulations.

Need a quotation, COA or samples of WELCYRO10?
Full product details on wellsunpharma.com · email anthony@wlsunshine.com

Request a Quote

This technical data sheet is for professional reference only. Specifications may be updated without notice; the COA supplied with each batch is the binding quality document. TDS-WELCYRO10 · Rev. 1 · Issued November 2026.