TECHNICAL DATA SHEET · ANTICOCCIDIALS

WELDECO60 – Decoquinate 6% Premix

Quinolone anticoccidial premix for in-feed use, containing 60 mg decoquinate per gram (6%).

Chickens | Mixed in feed | 25 kg/bag



TARGET ANIMALS

- Chickens (broilers)

WITHDRAWAL PERIOD

Chickens	5 days

INDICATIONS

- **Chickens:** prevention of coccidiosis caused by *Eimeria mitis*, *E. tenella*, *E. maxima*, *E. acervulina*, *E. necatrix* and *E. brunetti*.

Mode of action: Decoquinate mainly blocks the development of coccidial sporozoites, with peak activity on day 1 after infection. It suppresses immunity to coccidia, so broilers should be medicated continuously; rotate regularly to limit resistance.

DOSAGE & ADMINISTRATION

Animal	WELDECO60 per 1,000 kg feed	Duration
Broilers	453 g	7–14 consecutive days

Administration: mixed in feed. Dose expressed as the product; 453 g supplies about 27.2 g decoquinate per 1,000 kg feed.

ADVERSE REACTIONS

- No adverse reactions have been observed at the recommended dosage.

PRECAUTIONS

- Do not use in feed containing bentonite.

Please use all products in accordance with local laws and regulations.

Need a quotation, COA or samples of WELDECO60?
Full product details on wellsunpharma.com · email anthony@wlsunshine.com

Request a Quote

This technical data sheet is for professional reference only. Specifications may be updated without notice; the COA supplied with each batch is the binding quality document. TDS-WELDECO60 · Rev. 1 · Issued November 2026.