TECHNICAL DATA SHEET · ANTICOCCIDIALS

WELDICLA5 – Diclazuril 0.5% Premix

Triazine anticoccidial premix for in-feed use, containing 5 mg diclazuril per gram (0.5%).

Poultry | Rabbits | Mixed in feed | 25 kg/bag



TARGET ANIMALS

- Poultry
- Rabbits

WITHDRAWAL PERIOD

Chickens	5 days
Rabbits	14 days

INDICATIONS

- **Poultry and rabbits:** prevention of coccidiosis. Effective against chicken *Eimeria tenella*, *E. acervulina*, *E. necatrix*, *E. brunetti* and *E. maxima*, and against duck and rabbit coccidia.

Mode of action: Diclazuril is coccidiocidal and acts on all developmental stages, with peak activity on sporozoites and first-generation schizonts (the first 2 days of the life cycle).

DOSAGE & ADMINISTRATION

Animal	WELDICLA5 per 1,000 kg feed	Diclazuril in feed
Poultry	200 g	1 mg/kg
Rabbits	200 g	1 mg/kg

Administration: mixed evenly in feed. Dose expressed as the product.

ADVERSE REACTIONS

- No adverse reactions have been observed at the recommended dosage.

PRECAUTIONS

- Can be used in commercial feed and during the rearing period.
- Do not use in laying poultry producing eggs for human consumption during the laying period.
- Short duration of action: the effect drops markedly 1 day after stopping and is essentially gone after 2 days, so feed continuously.
- The inclusion rate is very low; mix thoroughly or efficacy will be affected.

Please use all products in accordance with local laws and regulations.

Need a quotation, COA or samples of WELDICLA5?
Full product details on wellsunpharma.com · email anthony@wlsunshine.com

Request a Quote

This technical data sheet is for professional reference only. Specifications may be updated without notice; the COA supplied with each batch is the binding quality document. TDS-WELDICLA5 · Rev. 1 · Issued November 2026.