TECHNICAL DATA SHEET · ANTICOCCIDIALS

WELDINI250 – Dinitolmide 25% Premix

Dinitro-class anticoccidial premix for in-feed use, containing 250 mg dinitolmide per gram (25%).

Chickens | Mixed in feed | 25 kg/bag



TARGET ANIMALS

- Chickens

WITHDRAWAL PERIOD

Chickens	3 days

INDICATIONS

- **Chickens:** coccidiosis. At the recommended dose it does not affect the development of immunity to coccidia, making it suitable for layer replacement and breeder flocks.

Mode of action: Dinitolmide has peak activity on day 3 after infection and also acts on oocyst sporulation.

DOSAGE & ADMINISTRATION

Animal	WELDINI250 per 1,000 kg feed	Dinitolmide per 1,000 kg feed
Chickens	500 g	125 g

Administration: mixed in feed. Dose expressed as the product.

ADVERSE REACTIONS

- No adverse reactions have been observed at the recommended dosage.

PRECAUTIONS

- Can be used in commercial feed and during the rearing period.
- Do not use in laying hens producing eggs for human consumption during the laying period.
- Withdrawing too early often leads to relapse, so broilers should be medicated continuously.
- Particle size affects efficacy; dinitolmide should be an ultra-fine powder.
- Above 250 mg dinitolmide per kg feed for more than 15 days, weight gain of chicks may be suppressed.

Please use all products in accordance with local laws and regulations.

Need a quotation, COA or samples of WELDINI250?
Full product details on wellsunpharma.com · email anthony@wlsunshine.com

Request a Quote

This technical data sheet is for professional reference only. Specifications may be updated without notice; the COA supplied with each batch is the binding quality document. TDS-WELDINI250 · Rev. 1 · Issued November 2026.