TECHNICAL DATA SHEET · ANTIBACTERIALS

WELFLOR50 SP – Florfenicol 5% Soluble Powder

Amphenicol antibiotic soluble powder for drinking water, containing 50 mg florfenicol per gram (5%).

Chickens | In drinking water | 1 kg foil bag



TARGET ANIMALS

- Chickens

WITHDRAWAL PERIOD

Chickens	5 days

INDICATIONS

- **Chickens:** treatment of bacterial diseases caused by florfenicol-susceptible bacteria.

DOSAGE & ADMINISTRATION

Animal	WELFLOR50 SP in drinking water	Florfenicol	Duration
Chickens	2–4 g per litre	100–200 mg per litre	3–5 days

Dose expressed as the product.

ADVERSE REACTIONS

- Strong immunosuppressive effect when used above the recommended dose.

PRECAUTIONS

- Do not use in laying hens producing eggs for human consumption during the laying period.
- Use with caution in breeder chickens.
- Do not use during vaccination or in animals with severely impaired immune function.
- Do not combine with macrolides or lincosamides (same 50S target, mutual antagonism); may antagonise penicillins or aminoglycosides.

Please use all products in accordance with local laws and regulations.

Need a quotation, COA or samples of WELFLOR50 SP?
Full product details on wellsunpharma.com · email anthony@wlsunshine.com

Request a Quote

This technical data sheet is for professional reference only. Specifications may be updated without notice; the COA supplied with each batch is the binding quality document. TDS-WELFLOR50-SP · Rev. 1 · Issued November 2026.