TECHNICAL DATA SHEET · ANTICOCCIDIALS

WELROBEN100 – Robenidine HCl 10% Premix

Guanidine anticoccidial premix for in-feed use, containing 100 mg robenidine hydrochloride per gram (10%).

Chickens | Rabbits | Mixed in feed | 25 kg/bag



TARGET ANIMALS

- Chickens
- Rabbits

WITHDRAWAL PERIOD

Chickens	5 days
Rabbits	7 days

INDICATIONS

- **Chickens and rabbits:** coccidiosis. Effective against chicken *Eimeria tenella*, *E. necatrix*, *E. brunetti*, *E. maxima*, *E. acervulina* and *E. mitis*, including strains resistant to other anticoccidials, and against rabbit coccidia.

Mode of action: Robenidine mainly inhibits first-generation schizonts and also acts on second-generation schizonts; peak activity is on day 3 after infection.

DOSAGE & ADMINISTRATION

Animal	WELROBEN100 per 1,000 kg feed	Robenidine HCl per 1,000 kg feed
Chickens	300–600 g	30–60 g
Rabbits	1,000–1,500 g	100–150 g

Administration: mixed in feed. Dose expressed as the product.

ADVERSE REACTIONS

- No adverse reactions have been observed at the recommended dosage.

PRECAUTIONS

- Can be used in commercial feed and during the rearing period.
- Do not use in laying hens producing eggs for human consumption during the laying period.
- Long-term or high-level feeding (60 mg/kg feed) can give meat and eggs an off-flavour; below 30 mg/kg feed this does not occur.
- Stopping too early often leads to relapse; medicate continuously.

Please use all products in accordance with local laws and regulations.

Need a quotation, COA or samples of WELROBEN100?

Full product details on wellsunpharma.com · email anthony@wlsunshine.com

Request a Quote

This technical data sheet is for professional reference only. Specifications may be updated without notice; the COA supplied with each batch is the binding quality document. TDS-WELROBEN100 · Rev. 1 · Issued November 2026.