TECHNICAL DATA SHEET · ANTIBACTERIALS

WELTIAMU100 – Tiamulin Fumarate 10% Premix

Pleuromutilin antibiotic premix for in-feed use, containing 100 mg tiamulin fumarate per gram (10%).

Pigs | Mixed in feed | 25 kg/bag



TARGET ANIMALS

- Pigs

WITHDRAWAL PERIOD

Pigs	7 days

INDICATIONS

- **Pigs:** prevention and treatment of mycoplasmal pneumonia and *Actinobacillus* pleuropneumonia, and swine dysentery caused by *Brachyspira* (treponema).

DOSAGE & ADMINISTRATION

Animal	WELTIAMU100 per 1,000 kg feed	Duration
Pigs	400–1,000 g	5–10 days

Administration: mixed in feed. Dose expressed as the product.

ADVERSE REACTIONS

- Skin erythema is occasionally seen at the normal dose. Overdose can cause transient salivation, vomiting and CNS depression.

PRECAUTIONS

- Do not use together with polyether ionophores such as monensin, salinomycin or narasin; this can cause growth depression, ataxia, paralysis and death.
- May reduce the efficacy of macrolides or lincomycin (same 50S binding site).
- Users should avoid contact with eyes and skin.
- Above 40 °C ambient temperature, medicated feed should not be stored for more than 7 days.

Please use all products in accordance with local laws and regulations.

Need a quotation, COA or samples of WELTIAMU100?
Full product details on wellsunpharma.com · email anthony@wlsunshine.com

Request a Quote

This technical data sheet is for professional reference only. Specifications may be updated without notice; the COA supplied with each batch is the binding quality document. TDS-WELTIAMU100 · Rev. 1 · Issued November 2026.